

FORM TP 2007247

TEST CODE **02134020** MAY/JUNE 2007

CARIBBEAN EXAMINATIONS COUNCIL ADVANCED PROFICIENCY EXAMINATION

PURE MATHEMATICS

UNIT 1 – PAPER 02

2 hours

(23 MAY 2007 (p.m.))

This examination paper consists of THREE sections: Module 1, Module 2 and Module 3.

Each section consists of 2 questions. The maximum mark for each section is 40. The maximum mark for this examination is 120. This examination consists of 6 pages.

INSTRUCTIONS TO CANDIDATES

1. **DO NOT** open this examination paper until instructed to do so.

2. Answer ALL questions from the THREE sections.

3. Unless otherwise stated in the question, any numerical answer that is not exact **MUST** be written correct to three significant figures.

Examination Materials

Mathematical formulae and tables Electronic calculator Graph paper

> Copyright © 2006 Caribbean Examinations Council® All rights reserved.

Section A (Module 1)

Answer BOTH questions.

(i) all the real factors of
$$g(x)$$
 [3 marks]
(ii) all the real roots of $g(x) = 0$.
(i) The function f is defined by $f(x) = x^4 - 9x^3 + 28x^2 - 36x + 16, x \in \mathbb{R}$; and
 $u = x + \frac{4}{x}, x \neq 0$.
(i) Express u^2 in terms of x .
(ii) By writing $f(x) = x^2 \left[x^2 - 9x + 28 - \frac{36}{x} + \frac{16}{x^2} \right]$ and using the result from
(b) (i) above, show that if $f(x) = 0$, then $u^2 - 9u + 20 = 0$.
(iii) Hence, determine the values of $x \in \mathbb{R}$ for which $f(x) = 0$.
(i) Let $S_n = \sum_{r=1}^n r$ for $n \in \mathbb{N}$. Find the value of n for which $3S_{2n} = 11 S_n$.
(a) Let $S_n = \sum_{r=1}^n r = \frac{1}{2}n(n+1)$]
(b) The quadratic equation $x^2 - nx + 24 = 0$, $n \in \mathbb{R}$ has roots g and β and the quadratic

(b) β , and the quadratic The quadratic equation $x^2 - px + 24 = 0$, $p \in \mathbb{R}$, has roots α a equation $x^2 - 8x + q = 0$, $q \in \mathbb{R}$, has roots $2\alpha + \beta$ and $2\alpha - \beta$.

(i)	Express p and q in terms of α and β .		[2 marks]	
(ii)	Find the values of α and β .		[4 marks]	
(iii)	Hence, determine the values of p and q .		[2 marks]	
Prove, by Mathematical Induction, that $n^2 > 2n$ for all integers $n \ge 3$.				
Total 20 marks				

GO ON TO THE NEXT PAGE

1.

(a)

Let $g(x) = x^4 - 9, x \in \mathbf{R}$. Find

- marks]
- mark]
- (b
 - marks]
 - ult from marks]
 - marks]

marks

2.

(c)

Section B (Module 2)

Answer BOTH questions.

3.

The circle shown in the diagram below (not drawn to scale) has centre C at (5, -4) and touches the y-axis at the point D. The circle cuts the x-axis at points A and B. The tangent at B cuts the y-axis at the point P.



Determine

(a)

(b)

(i)	the length of the radius of the circle	[2 marks]
(ii)	the equation of the circle	[1 mark]
(iii)	the coordinates of the points A and B , at which the cir	cle cuts the <i>x</i> -axis [6 marks]
(iv)	the equation of the tangent at B	[4 marks]
(v)	the coordinates of <i>P</i> .	[2 marks]
Show by calculation that $PD = PB$. [5 marks]		
		Total 20 marks

GO ON TO THE NEXT PAGE

4.

(a)

- Prove that $\cos 2\theta \equiv \frac{1 \tan^2 \theta}{1 + \tan^2 \theta}$. [4 marks] (i)
- Hence, show, without using calculators, that $\tan 67\frac{1}{2}^{\circ} = 1 + \sqrt{2}$. [**7 marks**] (ii)
- In the triangle shown below, (not drawn to scale), $\sin q = \frac{3}{5}$ and $\cos p = \frac{5}{13}$. (b)



Determine the exact values of

[1 mark] (i) $\cos q$ [1 mark] (ii) $\sin p$ sin r [3 marks] (iii) $\cos(p+t).$ [4 marks] (iv)

Total 20 marks

Section C (Module 3)

Answer BOTH questions.

Given that
$$y = \sqrt{5x^2 + 3}$$
,

5.

(a)

- (i) obtain $\frac{dy}{dx}$ [4 marks]
- (ii) show that $y \frac{dy}{dx} = 5x$ [2 marks]

(iii) hence, or otherwise, show that $y \frac{d^2 y}{dx^2} + \left(\frac{dy}{dx}\right)^2 = 5$. [4 marks]

(b) At a certain port, high tides and low tides occur daily. Suppose t minutes after high tide, the height, h metres, of the tide above a fixed point is given by

$$h = 2\left(1 + \cos\frac{\pi t}{450}\right), \ 0 \le t.$$

[Note: High tide occurs when h has its maximum value and low tide when h has its minimum value.]

Determine

- (i) the height of the tide when high tide occurs for the first time [2 marks]
- (ii) the length of time which elapses between the first high tide and the first low tide [3 marks]
- (iii) the rate, in metres per minute, at which the tide is falling 75 minutes after high tide. [5 marks]

Total 20 marks

GO ON TO THE NEXT PAGE

(i) Use the result
$$\int_{0}^{a} f(x)dx = \int_{0}^{a} f(a-x) dx, a > 0$$
, to show that if $I = \int_{0}^{\frac{\pi}{2}} \sin^{2}x dx$,

then
$$I = \int_{0}^{\pi/2} \cos^2 x \, dx$$
. [2 marks]

Hence, or otherwise, show that
$$I = \frac{\pi}{4}$$
.

[6 marks]

[4 marks]

(b)

(ii)

(i)

(a)

6.

- Sketch the curve $y = x^2 + 4$.
- Calculate the volume created by rotating the plane figure bounded by x = 0, (ii) y = 4, y = 5 and the curve $y = x^2 + 4$ through 360° about the y-axis.

[8 marks]

Total 20 marks

END OF TEST

02134020/CAPE 2007

a